Southern Cross University ePublications@SCU

Southern Cross Plant Science

2008

Evaluation of starch granule morphology in wild relatives of rice using scanning electron microscopy

Shabana Kasem
Southern Cross University

Daniel LE Waters
Southern Cross University

Nicole F. Rice Southern Cross University

Robert J. Henry
Southern Cross University

Publication details

Kasem, S, Waters, DLE, Rice, NF & Henry, RJ 2008, 'Evaluation of starch granule morphology in wild relatives of rice using scanning electron microscopy', paper presented to 58th Australian Cereal Chemistry Conference, Gold Coast Qld., 31 August – 4 September.

ePublications@SCU is an electronic repository administered by Southern Cross University Library. Its goal is to capture and preserve the intellectual output of Southern Cross University authors and researchers, and to increase visibility and impact through open access to researchers around the world. For further information please contact epubs@scu.edu.au.

Evaluation of starch granule morphology in wild relatives of rice using scanning electron microscopy


S. Kasem^{1*}, D.L.E. Waters², N. Rice1,² and R.J. Henry1,²

¹Grain Foods CRC, Southern Cross University PO Box 157 Lismore NSW 2480, ²Centre for Plant Conservation Genetics, PO Box 157 Lismore NSW 2480

Australia possesses a diverse range of wild rice species. Investigation of their morphology and starch properties may provide a platform for developing new hybrids or cultivars of cultivated rice. Starch functionality is influenced by ultra structure of endosperm and starch granule morphology^{1,2}. A scanning electron microscopy study was used to investigate the endosperm and starch ranule morphology in species of Australian wild rice.

Figure 2: Scanning Electron Microscopy images showing the different grain morphologies present between cultivated (Column A) and wild Oryza (Column B) species. P-pericarp; CG- compound granule; AL- aleurone layer; E-endosperm; PB- protein bodies; CW- cell wall; SG- starch granule.

Six seeds per species were manually de-husked and then fixed in formalin. Seeds then underwent a standard ethanol dehydration, were critical point dried and gold coated before being observed using a Leo440Stereoscan scanning electron microscope.



Poaceae

Figure 3: Grain colour and shape variation observed between the Erhartoideae. Images not to scale

Wild rice species showed both differences and similarities in their endosperm morphology when compared to cultivated rice. Species differences in cell thickness of the aleurone layers were observed. The frequency of protein bodies also differed amongst the wild species and between wild and cultivated species. Further investigation by physiochemical testing will reveal more information about the starch properties of wild species of rice.

The authors would like to thank Maxine Dawes from Environmental Science and Technology, Southern Cross University, for her technical assistance and Dr Frances Shapter for her technical advice. Also Dr Sally Dillon from Australian Tropical Crops and Forages Collection, Dept. Primary Industries and Fisheries, LMB 1 Biloela, QLD 4715. www. dpi.qld.gov.au/auspgris for supplying seed and Grain Foods CRC for funding this study.

References

- 1. Singh, N. and L. Kaur, Morphological, thermal, rheological and retrogradation properties of potato starch fractions varying in granule size. Journal of the Science of Food and Agriculture, 2004. 84: p. 1241-1252.
- 2. Londo, J.P., et al., Phylogeography of Asian wild rice, Oryza rufipogon, reveals multiple independent domestications of cultivated rice, Oryza sativa. Proc Natl Acad Sci, 2006. 103(25): p. 9578-83.

